

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/869, 5/06, A61K 39/245, 48/00, 35/76, C07K 14/035, C12N 15/38, A61P 35/00		A1	(11) International Publication Number: WO 00/65078 (43) International Publication Date: 2 November 2000 (02.11.00)		
(21) International Application Number: PCT/US00/11031		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).			
(22) International Filing Date: 24 April 2000 (24.04.00)					
(30) Priority Data: 09/299,817 26 April 1999 (26.04.99) US					
<p>(71) Applicant: CEDARS-SINAI MEDICAL CENTER [US/US]; 8700 Beverly Boulevard, Los Angeles, CA 90048 (US).</p> <p>(72) Inventors: WESCHLER, Steven, L.; 4401 Sevenoaks Court, Westlake Village, CA 92361 (US). NESBURN, Anthony, B.; 18128 Wakecrest Drive, Malibu, CA 90265 (US). PERN, Guey-Chuen; Apartment A, 217 South Fifth, Alhambra, CA 91801 (US). YU, John, S.; 1929 Linda Flora Drive, Los Angeles, CA 90077 (US). BLACK, Keith, L.; 1233 Roberto Lane, Los Angeles, CA 90077 (US).</p> <p>(74) Agents: STEINBERG, Nisan, A. et al.; Pretty, Schroeder & Poplawski, 19th Floor, 444 South Flower Street, Los Angeles, CA 90071 (US).</p>					
<p>(54) Title: A HERPES SIMPLEX VIRUS TYPE 1 (HSV-1)-DERIVED VECTOR FOR SELECTIVELY INHIBITING MALIGNANT CELLS AND METHODS FOR ITS USE TO TREAT CANCERS AND TO EXPRESS DESIRED TRAITS IN MALIGNANT AND NON-MALIGNANT MAMMALIAN CELLS</p> <p>(57) Abstract</p> <p>Disclosed is a method of selectively inhibiting the growth of malignant cells in mammals, including humans. The method selectively inhibits the growth of malignant cells of all varieties, and is particularly useful in treating brain tumors and other malignancies of the central nervous system. The method employs HSV-1-derived vectors containing a DNA having a deletion in both copies of the LAT gene and both copies of the ICP34.5 gene of HSV-1. The vectors are delivered to malignant cells either <i>in vivo</i> or <i>in vitro</i>, in accordance with the method. The HSV-1-derived expression vectors are non-neurovirulent and do not spontaneously reactivate from latency, and they optionally contain a functional HSV thymidine kinase gene, which can enhance the effectiveness against cancer of drug treatment with gancyclovir or acyclovir. Alternatively, the HSV-1-derived vectors contain at least one transcriptional unit of a LAT promoter sequence operatively linked to a nucleic acid having a nucleotide sequence encoding a polypeptide toxic for cells expressing the vector, for example, human interferon-γ. A method of expressing in a mammalian cell a gene encoding a preselected protein, a method of treating a genetic defect, and a method of detecting an HSV-1 expressing cell also employ vectors of the present invention that contain at least one transcriptional unit of a constitutive LAT promoter operatively linked to and controlling the transcription of a gene encoding a preselected protein. Also, disclosed are kits for expressing in a mammalian cell a gene encoding a preselected protein, useful for practicing the methods, and mammalian cells containing the HSV-derived vectors.</p>					

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**A HERPES SIMPLEX VIRUS TYPE 1 (HSV-1)-DERIVED VECTOR FOR
SELECTIVELY INHIBITING MALIGNANT CELLS
AND METHODS FOR ITS USE TO TREAT CANCERS AND
TO EXPRESS DESIRED TRAITS IN MALIGNANT AND NON-MALIGNANT
MAMMALIAN CELLS**

5

The U.S. Government has a paid-up license in this invention and the right in limited circumstances to require the patent owner to license others on reasonable terms as provided for by the terms of grants EY07566 and EY10243 awarded by the NIH Public Health Service.

BACKGROUND OF THE INVENTION

10 Throughout this application various publications are referenced within parentheses. The disclosures of these publications in their entireties are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains.

15 1. FIELD OF THE INVENTION

The present invention is related to the medical arts, particularly to the field of gene therapy.

20 2. DISCUSSION OF THE RELATED ART

Viruses have been tested for their ability to treat various types of malignancies in animals and humans. The proposed therapeutic mechanisms of viral tumor therapy in the prior art include: (i) 25 producing new antigens on the tumor cell surface to induce immunologic rejection, called "xenogenization," and (ii) direct killing of the tumor cell by a virus, called "oncolysis."

Several animal models and animal types of malignant tumor have been used to study oncolysis with wild-type viruses. (Moore, *Ann. Rev. Microbiol.* 8: 393 [1954]; Moore, *Progr. Exp. Tumor Res.* 1: 411 [1960]). At least nine viruses have been shown to be capable of inducing some degree of tumor regression of a variety of tumors in animals. A major drawback found in these early studies, however, was systemic infection of the patient by the virus.

30 Later in the quest for a viral therapy for cancer, clinical trials employing herpes viral vector therapy were approved in the United States to treat human tumors. (Culver, *Clin. Chem.* 40: 510 [1994]). These studies employed replication-incompetent or defective viruses to potentially overcome the problem of systemic infection by the virus. However, the use of replication-defective herpes viruses for treating malignant tumors requires producer cells because each replication-defective herpes virus particle can enter only a single cell and cannot productively infect others thereafter.

Thus, due to their inability to replicate, replication-defective herpes viruses cannot spread to other tumor cells, they are unable to penetrate a deep, multilayered tumor *in vivo*. (Markert *et al.*, *Neurosurg.* 77: 590 [1992]).

On the other hand, the herpes simplex virus type 1 (HSV-1) appears to be particularly well suited for use in the treatment of malignancies. Mutation of several of the viral genes involved in DNA replication, including UTPase and thymidine kinase- render the virus replication-defective in normal postmitotic cells, like neurons, but replication-competent in dividing cells, which can complement the defect. In addition, the HSV-thymidine kinase (TK) gene product can convert the anti-herpes drugs gancyclovir or acyclovir to nucleotide analogs which block both viral and cellular replication, thereby killing dividing tumor cells.

The need for a safe and effective HSV-1-derived vector is especially acute with respect to malignant tumors of the central nervous system. These malignancies are usually fatal, despite recent advances in the areas of neurosurgical techniques, chemotherapy and radiotherapy. In particular, there are no standard therapeutic modalities that can substantially alter the prognosis for patients with malignant tumors of the brain, cranium, and spinal cord. For example, high mortality rates persist for patients diagnosed with malignant medulloblastomas, malignant meningiomas, malignant neurofibrosarcomas and malignant gliomas, which are characterized by infiltrative tumor cells throughout the brain. Although intracranial tumor masses can be debulked surgically, treated with palliative radiation therapy and chemotherapy, the survival associated with a diagnosis of glioma, especially glioblastoma, is typically measured in months. Oldfield *et al.* introduced viral vectors carrying the herpes simplex virus (HSV)-1 thymidine kinase (HS-tk) gene into brain tumor cells in human patients. (Oldfield *et al.*, *Human Gene Therapy* 4: 39 [1993]). In this study, there was some evidence of anti-tumor effect in five of the eight patients in the clinical trial. However, none of the patients was cured of brain cancer. Some of the limitations of current viral based therapies, described by Oldfield, include: (1) the low titer of virus produced; (2) virus spread limited to the region surrounding the producer cell implant; (3) possible immune response to the producer cell line; (4) possible insertional mutagenesis and transformation of virally infected cells; (5) a single treatment regimen of the drug, gancyclovir, because the "suicide" product kills virally infected cells and producer cells; and (6) the bystander effect of killing being limited to cells in direct contact with the virally transformed cells. (Bi, W. L. *et al.*, *Human Gene Therapy* 4: 725 [1993]).

During the early 1990's, the use of genetically engineered replication-competent HSV-1 viral vectors was first explored in the context of finding an antitumor viral therapy. Replication-competent mutants of herpes simplex virus type I (HSV-1) with single mutations demonstrated therapeutic potential against experimental malignant brain tumors, while being attenuated for neurovirulence. It was thought that a replication-competent virus would have the advantage of being able to enter one tumor cell, make multiple copies of its genome, lyse the cell and spread to other tumor cells. A

thymidine kinase-deficient (TK⁻) mutant, *dlsptk*, was able to destroy human malignant glioma cells implanted into the brain of an animal. (Martuza *et al.*, *Science* 252: 854 [1991]). The major disadvantage to this system was that these TK⁻ mutants were only moderately attenuated for neurovirulence, i.e., the ability to replicate in brain cells causing inflammation of the brain, and they 5 produced encephalitis at the doses required to kill the tumor cells adequately. (J.M. Markert *et al.*, *Neurosurgery* 32: 597 [1993]). Roizman described a HSV-based TK⁻ vector system capable of expressing foreign genes inserted into the TK gene. (Roizman, *Herpes Simplex Virus as a Vector*, U.S. Patent Nos. 5,599,691 and 5,288,641). Residual neurovirulence of TK⁻ limits the usefulness of such vectors for tumor therapy.

10 Other single mutants of HSV-1 included hrR3, containing an insertion of the *Escherichia coli lacZ* gene into the viral ICP6 gene, which encodes the ribonucleotide reductase large subunit (T. Mineta *et al.*, *Gene Therapy* 1:S78 [1994]; T. Mineta *et al.*, *J. Neurosurg.* 80:381 [1994]) and R3616 containing deletions in both copies of the γ 34.5 gene, a neurovirulence gene. (Markert *et al.*, *Neurosurgery* 32:597 [1993]). Roizman described a recombinant, purportedly avirulent HSV lacking 15 the ability to express a functional γ 34.5 gene product, a neurovirulence factor. (Roizman, *Recombinant Herpes Simplex Viruses vaccines and methods*, U.S. Patent No. 5,328,688). Spontaneous reactivation rates of these mutants was only relatively attenuated, not entirely eliminated. (E.g., G.-C. Perng *et al.*, *J. Virol.* 70(5):2883-93 [1996]; G.-C. Perng *et al.*, *J. Virol.* 69(5):3033-411 [1995]).

20 Multi-mutated HSV-1 mutants have been described having augmented safety. Multiple mutations engineered into HSV-1 made the possibility of reversions of wild type unlikely and confirmed multiple and potentially synergistic safety features by attenuating of multiple mechanisms of virulence. Kramm *et al.* reported a HSV-1 mutant vector, MGH-1, defective for both ribonucleotide reductase and γ 34.5, which had higher therapeutic safety than hrR3, but had clearly 25 decreased therapeutic efficiency compared to hrR3. (C.M. Kramm *et al.*, *Therapeutic efficiency and safety of a second-generation replication-conditional HSV1 vector for brain tumor gene therapy*, *Hum. Gene Ther.* 8(17):2057-68 [1997]). Martuza *et al.* taught a replication competent HSV-1 vector having deletions in both of its γ 34.5 genes, as well as in the ICP6 gene which encodes for the large sub unit of the HSV ribonucleotide reductase. (Martuza *et al.*, *Replication-competent Herpes 30 Simplex Virus mediates destruction of neoplastic cells*, U.S. Patent No. 5,585,096). Martuza *et al.* taught a HSV-1-derived replication competent vector for tumor therapy that was driven by a tumor-specific or cell-specific promoter and was purportedly not neurovirulent. (Martuza *et al.*, *Tumor- or cell-specific Herpes Simplex Virus replication*, U.S. Patent No. 5,728,379). Although improved in safety, these mutants have shown attenuated therapeutic potential.

35 Even while some of these mutants, for example, γ 34.5 mutants, have been shown to reduce neurovirulence by as much as 100,000 fold in mice, a major problem with existing technology is the

spontaneous reactivation of HSV-1. Following infection at a peripheral site, HSV-1 establishes a life long latent infection of the sensory neurons enervating the peripheral site. In the absence of viral reactivation, latency is totally benign, with no known pathology. In rabbits, which can have rates of spontaneous HSV-1 reactivation comparable to those seen in humans, HSV-1 γ 34.5 mutants have greatly reduced spontaneous reactivation when rabbits are infected at “normal” doses (2×10^5 pfu/eye). (G.-C. Perng *et al.*, *An Avirulent ICP34.5 deletion mutant of Herpes Simplex Virus Type 1 is capable of in vivo spontaneous reactivation*, J. Virol. 69(5):3033-41 [1995]). In contrast, at extremely high infectious doses (2×10^8 pfu/eye) wild type spontaneous reactivation rates are achieved, separably from level of neurovirulence. (G.C. Perng *et al.*, *High-dose ocular infection with a Herpes Simplex Virus Type 1 ICP34.5 deletion mutant produces no corneal disease or neurovirulence yet results in wild-type levels of spontaneous reactivation*, J. Virol. 70(5):2883-93 [1996]).

During neuronal latency, latency-associated transcript (LAT) is the only viral gene that is abundantly transcribed. This is a function of the LAT promoter, a very powerful promoter with significant neuronal specificity. LAT is essential for efficient spontaneous reactivation. In rabbits, HSV-1 LAT null mutants have spontaneous reactivation rates of 1/3 or less than that of wild type or rescued viruses. (G.C. Perng *et al.*, *The Latency-Associated Transcript gene of Herpes Simplex Virus Type 1 (HSV-1) is required for efficient in vivo spontaneous reactivation of HSV-1 from latency*, J. Virol. 68(12):8045-55 [1994]). But this is still a substantial level of spontaneous reactivation.

Therefore, there is a desideratum for an effective HSV vector for brain tumor therapy that is non-neurovirulent and also does not reactivate spontaneously.

Delivery of the virus particles to tumor cells is another problem encountered in HSV therapy against brain tumors. Stereotactic inoculation of virus particles directly into a brain tumor has commonly been limited to small tumors, because the small volume of distribution by stereotactic inoculation limits the efficacy of viral therapy for large and disseminated tumors. (L.L. Muldoon *et al.*, *Comparison of intracerebral inoculation and osmotic blood-brain disruption for delivery of adenovirus, herpesvirus, and iron oxide particles to normal rat brain*, Am. J. Pathol. 147(6):1840-51 [1995]).

Kramm *et al.* described intrathecal injection of HSV vector hrR3 into the cerebrospinal fluid of rats with gliosarcomas, which resulted in HSV expression in frontal tumors and leptomeningeal tumor foci along the entire neuroaxis; however there was substantial toxicity associated with intrathecal injection of the vector. (C.M. Kramm *et al.*, *Herpes vector-mediated delivery of marker genes to disseminated central nervous system tumors*, Hum. Gene Ther. 7(3):291-300 [1996]).

Transvascular delivery of viral particles to tumor cells is hampered by the blood-brain barrier, and particularly the blood-tumor barrier, that results from the interendothelial tight junctions formed by cerebrovascular endothelial cells.

Neuwelt *et al.* used an intracarotid injection of hypertonic mannitol to osmotically disrupt the

blood-brain barrier. They reported that this enhanced the uptake by brain tissue of inactivated HSV-1 particles that were administered immediately afterward by bolus intracarotid injection, but there was no such enhancement when HSV was injected intravenously. (E.A. Neuwelt *et al.*, *Delivery of ultraviolet-inactivated 35S-herpesvirus across an osmotically modified blood-brain barrier*, *J. Neurosurg.* 74(3):475-79 [1991]; Also, S.E. Doran *et al.*, *Gene expression from recombinant viral vectors in the central nervous system after blood-brain barrier disruption*, *Neurosurgery* 36(5):965-70 [1995]; G. Nilaver *et al.*, *Delivery of herpesvirus and adenovirus to nude rat intracerebral tumors after osmotic blood-brain barrier disruption*, *Proc. Natl. Acad. Sci. USA* 92(21):9829-33 [1995]).

Rainov *et al.* described enhanced transvascular delivery of HSV into gliosarcoma cells in the brains of rats by injecting bradykinin, a vasoactive polypeptide, into the internal carotid artery to disrupt the blood-brain barrier. (N.G. Rainov, *Selective uptake of viral and monocrystalline particles delivered intra-arterially to experimental brain neoplasms*, *Hum. Gene. Ther.* 6(12):1543-52 [1995]; N.G. Rainov *et al.*, *Long-term survival in a rodent brain tumor model by bradykinin-enhanced intra-arterial delivery of a therapeutic herpes simplex virus vector*, *Cancer Gene Ther.* 5(3):158-62 [1998]).

A bradykinin analog, RMP-7, was shown to selectively open the blood-tumor barrier in rats to hrR3 HSV particles without the hypotensive effects associated with the use of bradykinin. (F.H. Barnett *et al.*, *Selective delivery of herpes virus vectors to experimental brain tumors using RMP-7*, *Cancer Gene Ther.* 6(1):14-20 [1999]).

There is a definite need for an effective HSV-1-derived vector for cancer therapy, that is virulence impaired and also does not reactivate spontaneously. There is a further definite need for a method of inhibiting malignant cells using such an HSV-1-derived vector that is also capable of reaching tumors of the central nervous system, particularly brain tumors. These and other benefits the present invention provides as described herein.

25 SUMMARY OF THE INVENTION

The present invention is directed to a method of selectively inhibiting the growth of malignant cells in mammals, including humans. The method selectively inhibits the growth of malignant cells of all varieties, and is particularly useful in treating brain tumors and other malignancies of the central nervous system. The method employs an HSV-1-derived viral vector containing a deletion in both 30 copies of the LAT gene and both copies of the ICP34.5 gene of HSV-1, and which is delivered to the malignant cell. One of the advantages of using a combined LAT null and γ 34.5 null HSV-1-derived vector, compared to LAT null mutants or γ 34.5 null mutants alone, is that regardless of the infectious dose, the double mutant does not reactivate from latency. The HSV-1-derived vector of the present invention employed in the method, optionally contains a functional HSV thymidine kinase gene, 35 which can enhance the effectiveness against cancer of drug treatment with gancyclovir or acyclovir.

The combination of LAT null and γ 34.5 null mutations results in a viral vector that does not damage neurons, but grows well and kills most other cells. Alternatively, the method employs an HSV-1-derived vector that contains at least one transcriptional unit of a LAT promoter sequence operatively linked to a nucleic acid having a nucleotide sequence encoding a polypeptide toxic for 5 cells expressing the vector, for example, human interferon- γ , interleukin-2, interleukin-4, interleukin-6, interleukin-10, interleukin-12, granulocyte-macrophage colony stimulating factor, tumor necrosis factor- α , Fas ligand, human connexin-43, VP-16, or VP-22.

The present invention relates to a method of expressing a preselected protein in a mammalian cell, employing a vector of the present invention that contains at least one transcriptional unit of a 10 constitutive LAT promoter operatively linked to a gene encoding any preselected protein. When the infected cell expresses, from the LAT promoter, it produces the preselected protein in an elevated amount compared to uninfected homologous cells. Thus, the present invention also relates to a method of treating a genetic defect in a mammal that is based on expressing a desired preselected protein, using a vector in accordance with the present invention, to mitigate or ameliorate a genetic 15 disease condition.

The present invention also relates to a method of detecting a mammalian cell that expresses HSV, employing an HSV-1-derived vector, with the same double deletion of LAT and ICP34.5 genes, and a fluorescent or light-emitting (green fluorescent protein) gene inserted into the vector under the transcriptional control of the LAT promoter. The resulting fluorescence or luminescence allows easy 20 detection of the location of the HSV-1 expressing cell, when tissue sections are examined under a fluorescent microscope or when cells are analyzed with flow-activated cell sorting (FACS).

The present invention is also directed to HSV-1-derived vectors and kits for expressing a preselected protein in a mammalian cell, useful for practicing the method, and to mammalian cells containing the HSV-derived vectors.

25 These and other advantages and features of the present invention will be described more fully in a detailed description of the preferred embodiments which follows.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a schematic of the structures of Δ LAT Δ 34.5 and Prom Δ LAT Δ 34.5-GFP compared to wild type HSV-1 genome.

30 Figure 2 illustrates the construction of Prom Δ LAT Δ 34.5-GFP.

Figure 3 illustrates cytotoxicity of Prom Δ LAT Δ 34.5-GFP in human U-87 MG glioma cells (Fig. 3A) and human glioblastoma multiforme primary cells (GBM-1; Fig. 3B, GBM-2; Fig. 3C).

Figure 4 shows human U-87 MG glioma cells (Fig. 4A) and human glioblastoma multiforme primary cells (GBM-1; Fig. 4B, GBM-2; Fig. 4C), under phase contrast (left panels) and with

fluorescent microscopy showing fluorescence secondary to expression of green fluorescent protein from Prom Δ LAT Δ 34.5-GFP (right panels).

Figure 5 shows non-malignant human normal neuronal HCN-1A cells (Fig. 5A), astrocyte HFA cells (Fig. 5B) and endothelial HUVEC cells (Fig. 5C) infected with wild type HSV-1 (center panels) or with Prom Δ LAT Δ 34.5-GFP (right panels) at a MOI of 0.1 pfu/cells. Control cells were mock-infected (left panels). Bar length is 100 μ m.

Figure 6 shows plaque formation as dose-dependent with respect to gancyclovir concentration in confluent monolayers of rabbit skin (RS) cells infected with 100 pfu of wild type HSV-1 or with Prom Δ LAT Δ 34.5-GFP.

Figure 7 shows a Kaplan-Meier survival analysis for nude mice with brain tumors treated with intratumoral injection of Prom Δ LAT Δ 34.5-GFP with or without subsequent treatment with gancyclovir (GCV).

Figure 8 shows brain sections from nude mice that were stereotactically implanted with malignant U-87 MG cells (Fig. 8A-E) and were later inoculated intratumorally with Prom Δ LAT Δ 34.5-GFP (Fig. 8B-E) and treated with gancyclovir (Fig. 8E only), or of controls (Fig. 8A). Bar length is 50 μ m.

Figure 9 shows histological sections of malignant U-87 MG cells from 100-day survivor mice treated with Prom Δ LAT Δ 34.5-GFP, with subsequent gancyclovir treatment (Fig. 9B) or without gancyclovir treatment. Left panels are HE stained slides; right panels are slides viewed with fluorescence microscopy to show fluorescence secondary to expression by cells of green fluorescent protein from Prom Δ LAT Δ 34.5-GFP. Bar length is 50 μ m.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is directed to a method of selectively inhibiting the growth of malignant cells in mammals, including humans, non-human primates, for example simians, rodents such as mice and rats, gerbils, hamsters, and rabbits. The method selectively inhibits the growth of malignant cells of all varieties, compared to the effect on non-malignant cells, in malignant tumors in any body tissue or cavity, including malignant tumors in the brain, spine, skull, thorax, lung, peritoneum, prostate, ovary, uterus, breast, stomach, liver, bowel, colon, rectum, bone, lymphatic system, and skin. The method is useful in the treatment of primary malignancies and metastases. The method is useful in inhibiting the growth of malignant cells, including, but not limited to malignant cells of gliomas, glioblastomas, oligodendrogiomas, astrocytomas, ependymomas, primitive neuroectodermal tumors, atypical meningiomas, malignant meningiomas, neuroblastomas, sarcomas, lymphomas, and carcinomas of all kinds, including hepatocarcinomas or adenocarcinomas.

In the inventive method, an HSV-1-derived viral vector containing in its DNA a deletion in both copies of the LAT gene and both copies of the ICP34.5 gene of HSV-1, is delivered to the

malignant cell. Both of these genes are located in the viral long repeat of the wild type HSV-1 genome and therefore, in wild type HSV-1 virus both genes are present in two copies. (Fig. 1A). One of the advantages of using a combined LAT null mutant and γ 34.5 null mutant HSV-1-derived vector, compared to LAT null mutants or γ 34.5 null mutants, is that regardless of the infectious dose, the 5 double mutant does not reactivate from latency. Transcripts encoding LAT gene product or ICP34.5 gene product are not detectably produced in a cell infected with the mutated virus. Thus, following the use of the method in inhibiting the growth of malignant cells in the brain or other tissue, or in delivery of foreign genes to malignant or non-malignant neurons or other cell types, there is no concern that spontaneous reactivation or virulence might occur at future times, with possible 10 clinically harmful effects. Unlike wild type HSV-1, acute infection with a combined LAT-deletion/34.5-deletion mutant vector produces no corneal disease or other pathology.

Most preferably, the HSV-1-derived viral vector that is delivered to a malignant cell in accordance with the present method is a viral vector derived from HSV-1 strain McKrae. Strain McKrae is much more vigorous and virulent than other HSV-1 strains previously used as potential 15 vectors, such as KOS, F, and 17syn+. It is an advantage of HSV-1-derived vectors derived from strain McKrae that they have the ability to efficiently kill cultured cells in vitro. This has not been demonstrated for other strains of HSV-1 virus.

Preferably, but not necessarily, the HSV-1-derived vector used in the method comprises a functional HSV thymidine kinase gene. When the HSV-1-derived viral vector is delivered to a 20 malignant cell and internalized, the expressing host cell expresses transcripts encoding HSV thymidine kinase. The HSV thymidine kinase that is thereby produced in the expressing malignant cell can convert the drugs gancyclovir or acyclovir to a nucleotide analog within the malignant cell, inhibiting its growth. Thus, when an HSV-1-derived viral vector is employed that contains a 25 functional HSV thymidine kinase gene, these drugs become particularly effective in treating malignancies.

The selectivity of the method lies in the replication conditional nature of the HSV-1-derived viral vector. HSV-1 can replicate and go into a lytic phase in actively growing cells, because the cell provides the metabolic background for HSV-replication. The HSV-1-derived vectors of the present invention are not selective in infecting cells. They infect non-malignant cells, such as normal neurons 30 and glial cells just as well as they infect malignant cells, but they can only enter a lytic phase in rapidly growing malignant cells. Since the viral replication is conditional only in growing malignant cells, more of these malignant cells are preferentially inhibited, compared to the general population of less metabolically active non-malignant cells.

In another preferred embodiment of the method, the HSV-1-derived viral vector is an 35 expression vector that further comprises at least one transcriptional unit of a LAT promoter sequence, or operative fragment thereof, operatively linked to a nucleic acid having a nucleotide sequence

encoding a desired preselected protein, the presence of which polypeptide or protein results in toxicity to a malignant cell. A preferred example of a protein toxic to a malignant cell is human interferon- γ , or a functional fragment or fusion protein derived therefrom. When a malignant cell expresses a human interferon- γ gene from the LAT promoter sequence, the malignant cell constitutively produces 5 higher levels of human interferon- γ , compared to homologous malignant cells not expressing the HSV-1-derived vector, which acts, like other cytokines, in a paracrine manner to produce a cytotoxic immune response in the vicinity where the cytokine is secreted, thus preferentially inhibiting or killing other malignant cells close by in a tumor mass.

Other preferred polypeptides or proteins toxic to malignant cells include, but are not limited 10 to, interleukin-2, interleukin-4, interleukin-6, interleukin-10, interleukin-12, granulocyte-macrophage colony stimulating factor, tumor necrosis factor- α , Fas ligand (e.g., of human or murine origin), human connexin-43, VP-16, VP-22, or functional fusion proteins derived from any of these.

Inhibition of the growth of malignant cells is determined by directly counting cells 15 microscopically or by measuring physiologic or metabolic activity of malignant cells, in comparison with before-treatment values. For example, cell growth can be measured by evaluating the incorporation of [3 H]-thymidine into DNA during division or other known metabolic assessment techniques. Alternatively, inhibition of the growth of malignant cells is determined by measuring changes in the size of tumors comprising the malignant cells.

The present invention also relates to a method of expressing a gene encoding a preselected 20 protein in a mammalian cell, whether a non-malignant or malignant cell. The method can be used to infect cells *in vivo*, i.e., within a mammal, or to infect cells grown *in vitro*. The method involves delivering to a mammalian cell an HSV-1-derived vector which contains a deletion in both copies of the LAT gene and both copies of the ICP34.5 gene of HSV-1, and contains at least one transcriptional 25 unit of the constitutive LAT promoter sequence, or operative fragment thereof, operatively linked to a nucleic acid having a nucleotide sequence encoding any preselected protein, or operative fragment thereof. The protein can be one that is normally expressed by homologous uninfected cells, or, optionally, it can be a foreign protein. The cell expresses, from the LAT promoter, the nucleic acid encoding the desired preselected protein, in an elevated amount compared to uninfected homologous cells. Thus, the method can be used to deliver a foreign gene to "cure" a genetic defect.

Consequently, the present invention is also related to a method to treat a genetic defect in 30 a mammal, i.e., a disease condition in a mammal that can be mitigated or ameliorated by expression of a specific exogenous gene, such as Alzheimer's disease, Multiple Sclerosis, Parkinson's disease, Gaucher's disease, or Huntingdon's disease in humans.

Many mammalian cell types have receptors for binding HSV-1, and binding of an HSV-1- 35 derived vector results in the vector being internalized into the cell. However, for particular applications, internalizing of the HSV-1-derived vector is accomplished using any suitable method.

Preferably, a bispecific antibody can be prepared to facilitate selective internalization of the HSV-derived vector into specific cell types. Such a bispecific antibody contains at least two covalently linked antibodies, one binding an HSV-1 specific epitope and another specifically binding an epitope specific for a particular cell type of interest. The bispecific antibody is first incubated with the HSV-derived vector before delivery, in order to allow the vector to preferentially infect a particular cell type.

The present invention also relates to a method of detecting the presence of a cell expressing HSV-1, which like the method of expressing a gene encoding a preselected protein in a mammalian cell, involves delivering to a mammalian cell an HSV-1-derived vector that contains a deletion in both 10 copies of the LAT gene and both copies of the ICP34.5 gene of HSV-1, and contains at least one transcriptional unit of a LAT promoter sequence, or operative fragment thereof, operatively linked to a reporter gene, particularly, a nucleic acid having a nucleotide sequence encoding a fluorescent or light-emitting protein, or an operative fragment thereof. Most preferably, the reporter gene, is a gene encoding Green Fluorescent Protein ([GFP]; or enhanced Green Fluorescent Protein [EGFP]), 15 Yellow Fluorescent Protein, Blue Fluorescent Protein, a phycobiliprotein, such as phycoerythrin or phycocyanin, or any other protein which fluoresces under light of suitable wave-lengths of light. The HSV-expressing cell is then detected via fluorescence therefrom.

Another preferred reporter gene suitable for some applications is a gene encoding a protein that can enzymatically lead to the emission of light from a substrate(s); for purposes of the present 20 invention, such a protein is a "light-emitting protein." For example, a light-emitting protein includes proteins such as luciferase or apoaequorin. The HSV-expressing cell is then detected via luminescence therefrom.

The cells are detected, sorted, isolated or selected from cells not expressing the HSV-1-derived expression vector with the aid of, for example, a flow-activated cell sorter (FACS) set at the 25 appropriate wavelength(s). Alternatively, they are detected, isolated or selected manually from non-HSV-expressing cells using conventional microscopic technology.

For purposes of the present invention, "operatively linked" means that the promoter sequence, is located upstream from the coding sequence and that both sequences are oriented in a 5' to 3' manner with respect to each other, such that together they form a transcriptional unit from which 30 transcription could take place in vitro in the presence of all essential enzymes, transcription factors, co-factors, activators, and reactants, under favorable physical conditions, e.g., suitable pH and temperature. This does not necessarily mean that, in any particular cell, conditions will favor transcription.

Delivering an HSV-1-derived vector to a mammalian cell can be by any suitable method. In 35 some applications directed to malignant cells, the HSV-1-derived vector is preferably delivered by intratumoral injection through a surgical incision, for example, through a craniotomy for a brain

tumor. Typically, surgical debulking of a tumor is done, if possible, before injection of the HSV-1-derived vector into the remaining tumor mass for infection of malignant cells.

Another preferred delivery method, useful for treating gliomatosis or carcinomatosis, is intrathecal injection of the vector. Another preferred delivery method is transvascular delivery, by 5 intravenous or intra-arterial injection of HSV-1-derived viral particles.

For treating brain tumors, intra-arterial injection into the carotid artery is most preferred. It is preferable to facilitate passage of the vector across the blood-brain barrier. This can be accomplished, for example, by first disrupting the blood-brain barrier with a hypertonic (typically, 10 25%) mannitol infusion, immediately before or substantially simultaneously with the intra-arterial injection of HSV-1-derived vector. Alternatively, a vasoactive agent, such as bradykinin, or a functional analog, for example, RMP-7 (typically, about 1.5-3.0 μ g/kg body weight over 10-15 minutes; Barnett et al. [1999]) is used to facilitate passage across the blood-brain barrier. Bradykinin and its analogs transiently and preferentially open the blood-tumor barrier in the neovasculature of brain tumors.

15 For treating a brain tumor, another preferred delivery method is stereotactic inoculation of the viral vector into the malignant tumor at a site having pre-established coordinates.

The foregoing are merely illustrative and not an exhaustive description of suitable methods for delivering the HSV-1-derived vector in accordance with the methods of the present invention.

20 The present invention also relates to an HSV-1-derived vector for use in any of the present methods. As described above, the HSV-1-derived vector contains in its DNA a deletion in both copies of the LAT gene and both copies of the ICP34.5 gene of HSV-1. A preferred vector is derived from HSV-1 strain McKrae.

A preferred HSV-1-derived vector for use in the method is a viral vector designated 25 Δ LAT Δ 34.5, or a derivative thereof. Δ LAT Δ 34.5 is derived from HSV-1 strain McKrae. The DNA structure of Δ LAT Δ 34.5 is shown in Figure 1B; both copies of γ 34.5 and LAT are deleted (Fig. 1B). The deletions include the promoters for both genes, the entire γ 34.5 structural gene, and the first 1.67 kb of the structural region of LAT. This region of LAT contains the LAT function that results in enhanced spontaneous reactivation. Δ LAT Δ 34.5 contains a fully functional HSV thymidine kinase 30 gene. The Δ LAT Δ 34.5 vector replicates preferentially in malignant cells and selectively kills or inhibits the growth of malignant cells.

The construction of Δ LAT Δ 34.5 is illustrated in Figure 1. The top of the figure shows the 35 prototypic organization of the wild type HSV-1 genome. The TR_L (terminal repeat long) and the IR_L (internal or inverted repeat long) each contain identical copies of the LAT region, including the gene for ICP34.5 that overlaps a portion of the LAT gene in the opposite orientation.

Figure 1A shows a blow up of the LAT region, identical in both long repeats. The primary LAT transcript is 8.3 kb long. The solid rectangle represents a very stable portion of the LAT transcript called the 2 kb LAT. The open rectangle indicates the location of the LAT promoter. TATA indicates the LAT TATA-box within the promoter. The small arrow (at +1) indicates the start 5 of LAT transcription. *EcoRV*, *StyI*, and *HpaI* are restriction sites that define boundaries of deleted areas shown below. The relative size and location of ICP34.5 is shown above LAT. The open rectangle indicates the ICP34.5 open reading frame. The location of ICP0 is shown for reference.

Figure 1B shows Δ LAT Δ 34.5, containing 2 deletions in the LAT region. The first deletion 10 extends from the *EcoRV* restriction site at LAT nucleotide -161 to the *HpaI* restriction site at LAT nucleotide 1,673. This deletion removes the core LAT promoter and the first 1.67 kb of the LAT transcript. The second deletion encompasses the region corresponding to LAT nucleotides 6.2-7.1 kb and thus all of the gene for ICP34.5. Both copies of the LAT/ICP34.5 region (1 in each long repeat) are identical. Δ LAT Δ 34.5 therefore makes no LAT RNA transcripts and no ICP34.5 RNA transcripts.

15 Another preferred example of the HSV-1-derived vector of the present invention is a viral vector designated Prom Δ LAT Δ 34.5 (Fig. 1C), or a derivative thereof. Prom Δ LAT Δ 34.5 is derived from HSV-1 strain McKrae, and is particularly derived from Δ LAT Δ 34.5, sharing its ability to selectively kill or inhibit the growth of malignant cells. Like Δ LAT Δ 34.5, Prom Δ LAT Δ 34.5 is a 20 LAT null mutant and a γ 34.5 null mutant, also having a functional HSV thymidine kinase gene. It contains a DNA having the LAT promoter, a *BamHI* site cloning site into which foreign (xenogeneic) genes, i.e., genes not naturally expressed from the LAT promoter, such as those encoding GFP or human interferon- γ , can be inserted. Prom Δ LAT Δ 34.5 also has a Poly A signal site. (Fig. 1 and 2). This vector is capable of long term high level expression of any foreign (xenogeneic) or exogenous gene in neurons or other cells.

25 In Prom Δ LAT Δ 34.5, the region of the LAT promoter from the *EcoRV* site to the start of the LAT transcript and the region of LAT from the start of LAT transcription to the *StyI* site at nucleotide 76 have been restored to Δ LAT Δ 34.5. The original *StyI* site at LAT nucleotide 76 and the end of the original deletion at the *HpaI* site at LAT nucleotide 1673 have been replaced by a single unique *BamHI* site into which a xenogeneic gene can be inserted and expressed under the transcriptional 30 control of the LAT promoter.

The construction of Prom Δ LAT Δ 34.5 is further illustrated in Figure 2. The plasmid pLAT1.5 was constructed; pLAT1.5 contains the LAT promoter from LAT nucleotides -799 to +76 (i.e., the first 76 nucleotides of the LAT transcript), followed by a small multiple cloning site, a Poly A signal site, a unique *BamHI* site, and LAT nucleotides 1,673 to 4,656. The foreign gene GFP was cloned 35 into the multiple cloning site to produce pLAT1.5-GFP, but any other foreign gene encoding a preselected protein can be cloned into the multiple cloning site instead. Cells were co-infected with

pLAT1.5-GFP and Δ LAT Δ 34.5 infectious DNA to produce Prom Δ LAT Δ 34.5-GFP by homologous recombination. The structure of the LAT region in the DNA of Prom Δ LAT Δ 34.5-GFP is thus as follows. The LAT promoter (including all upstream sequences) up to LAT nucleotide 76, followed by a foreign gene (GFP), followed by a Poly A site, followed by the remainder of LAT starting at 5 LAT nucleotide 1,673. The region of LAT from 77 to 1,672 is therefore deleted and no LAT can be made. The genomic location containing the gene for ICP34.5 (and therefore the corresponding of region of LAT from 6.2-7.1kb) is also deleted.

Prom Δ LAT Δ 34.5 can contain any foreign gene. The foreign gene is inserted into the cloning site of plasmid vector pLAT1.5. In Figure 2, GFP is used as an illustrative example of such a foreign 10 gene. The foreign gene can then be inserted into Δ LAT Δ 34.5 by homologous recombination of genomic Δ LAT Δ 34.5 DNA with the pLAT 1.5 plasmid vector DNA containing the foreign gene, as shown in Fig. 2 for the foreign gene GFP. In the final HSV-1-derived vector (in this case Prom Δ LAT Δ 34.5-GFP), the foreign gene is under the control of the LAT promoter and is followed by a Poly A site. The LAT promoter is highly active in a wide variety of cell types, and particularly 15 in neurons, producing long term expression of the foreign gene, even after the virus has become latent. This is a significant improvement over HSV-1 vectors using other viral promoters, since these other promoters are unlikely to provide long term expression.

The present invention also relates to a mammalian cell containing an HSV-1-derived vector 20 of the present invention, or unpackaged (unencapsidated) DNA thereof. The mammalian cell is a malignant cell or a non-malignant cell, originating in any mammalian species, but preferably derived from or contained in a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit. The cell is grown in vitro or is situated in vivo. A preferred malignant cell of the present invention is derived 25 from or is contained in a glioma, glioblastoma, oligodendrolioma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma. Examples of preferred non-malignant cell include neuronal, 30 epidermal, osteogenic, lymphatic, endothelial, epithelial, hepatic, breast, ovarian, uterine, or prostate cells, but a cell derived from any other kind of mammalian tissue is also preferred.

The present invention also relates to a kit for expressing in a mammalian cell a gene encoding 35 a preselected protein. The kit is a ready assemblage of materials for facilitating the practice of the present method or use of the present HSV-1-derived vector to infect a mammalian cell. A kit of the present invention contains at least an HSV-1-derived vector of the present invention and instructions for its effective use. The materials or components assembled in the kit are provided to the practitioner stored in any convenient and suitable way that preserves their operability and utility. For example the components can be in dissolved, dehydrated, or lyophilized form; they can be provided at room, refrigerated or frozen temperatures.

The present invention provides LAT null/ICP34.5 null HSV-1-derived vectors that are non-virulent and are not spontaneously reactivated. These vectors are safer and more effective than previously available HSV-derived vectors for the treatment of malignancies, and are particularly useful in the treatment of brain tumors *in vivo*. In addition, the HSV-1-derived vector of the present invention is able to inhibit the growth of malignant cells in *in vitro* culture.

5 The foregoing descriptions of the methods, vectors, cells and kits of the present invention are illustrative and by no means exhaustive. The invention will now be described in greater detail by reference to the following non-limiting examples.

EXAMPLES

10 **Example 1: Culture of Cells and HSV-1.**

Two kinds of human glioblastoma multiforme (GBM) primary cells were isolated from two different surgically obtained GBM specimens. The human glioblastoma cell line U87 MG, normal neuron HCN-1A cells, and human umbilical endothelial cells (HUVEC) were purchased from American Type Culture Collection (Rockville, MD). Normal human astrocyte HFA cells were kindly 15 provided by Carlo Tornatore (Laboratory of Molecular Medicine and Neuroscience, National Institute of Neurological Disorder and Stroke, Bethesda, MD). All cells except HCN-1A and RS cells were grown in Ham's F12/DME containing 10% fetal bovine serum and antibiotics. HCN-1A and rabbit skin (RS) cells were cultured in Dulbecco's minimum essential medium supplemented with 10% fetal bovine serum and antibiotics. Aseptic technique was used in maintaining all cell cultures. HSV-1 20 strain McKrae (wild type) and all HSV-1-derived expression vectors were routinely stored in tissue culture medium at -70°C until use.

Example 2: Determination of Cytotoxicity in In Vitro Cell Culture.

1x10³ cells of U 87MG or of the two GBM primary cell lines (GBM-1 and GBM-2) were seeded separately in each well of 96-well plates. Twenty four hours after seeding, PromΔLATΔ34.5-GFP was delivered to these cells at various titers. Three, seven and eleven days after infection with this viral vector, cell viability was determined using cell proliferation reagent WST-1 (Boehringer 25 Mannheim, Indianapolis, IN), which indicates the overall activity of mitochondrial dehydrogenases, and by measuring the absorbance of the dye solution in each well with a multi-well spectrophotometer. Viability of cells in the absence of PromΔLATΔ34.5-GFP represented 100 % 30 viability. As shown in Figure 3, the viable numbers of all three kinds of cells decreased with time in a dose-dependent manner in response to exposure to PromΔLATΔ34.5-GFP.

Example 3: In Vitro Cellular Expression of Green Fluorescent Protein (GFP) from Prom Δ LAT Δ 34.5-GFP.

To detect GFP expression from the LAT promoter, U 87MG and the two GBM primary cell lines (GBM-1 and GBM-2) were separately seeded in 4-well chamber slides (2×10^4 cells /well).

5 Twenty-four hours after seeding, Prom Δ LAT Δ 34.5-GFP was delivered at a MOI of 0.1 pfu/cells. On days 1, 2, 3 and 5, cells were fixed with 4 % paraformaldehyde in PBS for 30 min, washed, mounted and examined with fluorescence microscopy using a filter for GFP fluorescence.

10 Cellular expression of GFP after 3 days is shown in the left-hand panels of the micrographs in Figure 4. The right-hand panels show the same cells under phase contrast. In U87 MG cells, GFP-positive cells were barely detectable on day 1 post-infection. They showed GFP positivity and widespread cytopathic effects on day 3. (Fig. 4A). On day 5, almost all cells expressed GFP and a considerable number of cells had died and/or sloughed from the plate surface. In GBM-1, GFP-positivity was observed in a few cells on day 1 post-infection. On day 3, the vast majority of the cells were positive for GFP and exhibited cytopathic effects. (Fig. 4B). By day 5, almost all cells were 15 dead and/or sloughed from the plate surface. GBM-2 cells behaved in a manner similar to GBM-1. (Fig. 4C).

Example 4: Determinations of Infectivity.

20 To determine infectivities of wild type HSV-1 (strain McKrae) or Prom Δ LAT Δ 34.5-GFP with respect to non-malignant cells, neuronal (HCN-1A), astrocyte (HFA) and endothelial cells (HUVEC), were separately seeded in 24-well plates (2×10^4 cells per well). Twenty four hours after seeding, HCN-1A, HFA or HUVEC cells were infected with HSV-1 wild type strain McKrae or Prom Δ LAT Δ 34.5-GFP at a MOI of 0.1 pfu/cells. The cells were examined with a phase contrast microscope on days 3, 5 and 7. In each assay, controls were mock-infected.

25 Micrographs taken on day 3 are shown in Figure 5. On day 3, the majority of HCN-1A cells infected with wild type HSV-1 revealed morphological changes such as neuronal differentiation. (Figure 5A, center panel). Virtually all HCN-1A cells infected with Prom Δ LAT Δ 34.5-GFP remained normal (Fig. 5A, right panel). By day 5, all wild type HSV-1-infected HCN-1A cells had been killed. In contrast, only about a half of the cells infected with Prom Δ LAT Δ 34.5-GFP demonstrated a slight 30 morphological change.

30 Virtually all HFA cells infected with wild type HSV-1 were killed by day 3 (Fig. 5B, center panel), but HFA cells infected with Prom Δ LAT Δ 34.5-GFP showed few morphological changes. (Fig. 5B, right panel). On day 5, only a few HFA cells infected with Prom Δ LAT Δ 34.5-GFP showed any morphological change. HUVEC cells behaved in a manner similar to HFA cells. On day 3, about 90 % of HUVEC infected with wild type HSV-1 were killed (Fig. 5C, center panel), but all HUVEC cells 35 infected with Prom Δ LAT Δ 34.5-GFP were normal. (Fig. 5C, right panel). By day 5, there were no

viable HUVEC cells infected with wild type HSV-1. Cells infected with Prom Δ LAT Δ 34.5-GFP were all viable, but in some HUVEC cells a foamy structure appeared in the nucleus. Mock-infected control cells of each of the three cell lines showed no morphological changes. (Fig. 5A-C, left panels).

5

Example 5: In Vitro Sensitivity to Drug Treatment With Gancyclovir.

Confluent monolayers of RS cells in 6-well plates were infected with 100 pfu of wild type HSV-1 or infected with Prom Δ LAT Δ 34.5-GFP in 500 μ l of tissue culture medium for a 1- hour incubation period at 37°C. After the viral inoculum was removed, minimal essential medium (MEM) containing 5 % fetal bovine serum, 1 % methylcellulose and various concentrations of gancyclovir, were added to triplicate cultures and cells were incubated at 37°C for 3 days. Plaques were visualized by Giemsa stain and were counted.

Figure 6 shows gancyclovir sensitivities in RS cells hosting either wild type HSV-1 or Prom Δ LAT Δ 34.5-GFP. Plaque reduction assays demonstrated that cells infected with Prom Δ LAT Δ 34.5-GFP are as sensitive or more sensitive to gancyclovir, compared to cells infected with wild type HSV-1. (Fig. 6). The numbers of plaques in the wells without gancyclovir represents 100 %.

Example 6: Prom Δ LAT Δ 34.5-GFP in Mammalian Cells In Vivo.

Mammals. Six-week-old female mice (BALB/c) or nude mice (BALB/c-*nu/nu*) were used for all animal experiments and were obtained from Charles River (Wilmington, MA). Safety of Prom Δ LAT Δ 34.5-GFP In Vivo. To confirm the non-neurovirulence of Prom Δ LAT Δ 34.5-GFP, 18 mice (BALB/c) were stereotactically inoculated with 5 μ l of sterile culture medium containing either Prom Δ LAT Δ 34.5-GFP or wild type HSV-1. Mice were equally divided into 3 groups; Group A mice received 2 \times 10⁵ pfu wild type HSV-1; Group B mice received 2 \times 10⁵ pfu Prom Δ LAT Δ 34.5-GFP; and Group C received 1 \times 10⁷ pfu Prom Δ LAT Δ 34.5-GFP. Mice were followed for 60 days post-inoculation.

All 6 mice in Group A died within 6 days after infection. All 6 mice in Group B, inoculated with Prom Δ LAT Δ 34.5-GFP, were alive after 60 days. One mouse out of the six in Group C, died on day 9 post-inoculation.

Survival of Mammals with Malignancies After Delivery of Prom Δ LAT Δ 34.5-GFP and Responses to Gancyclovir Treatment. Survival studies were carried out to determine the efficacy of treatment with the present method of inhibiting the growth of a malignant cell. Animals were monitored and when they met criteria, they were euthanised. Criteria included moribund state, inability to feed, cachexia, and inability to support weight or to ambulate. If animals survived over 3 months, they were euthanised (via CO₂ asphyxiation) and their brains were harvested for histology.

For surgical procedures, all mice were anesthetized with an intraperitoneal injection of ketamine (100 mg/kg) and xylazine (10 mg/kg).

U87MG cells (2×10^5) were stereotactically implanted in the right basal ganglia of 30 nude mice. After anesthesia, a linear skin incision was made on the midline of the skull. A burr hole was 5 made on the coronal suture approximately 2 mm lateral to the midline on the right side. After the mice were immobilized in a stereotactic apparatus, 2×10^5 U87 MG cells in 2 μ l of Ham's F12/DME medium, containing 1.2 % methylcellulose, were injected with a Hamilton syringe to a depth of 2 mm from the cortical surface over a period of 2 min. Before retracting, the needle was left as it was for 2 additional min. The skin incision was closed using sterile surgical staples. Thereafter, animals 10 were divided into 3 groups of ten mice each.

Seven days after tumor cells were implanted, the first group of mice received stereotactic inoculation of 4.4×10^6 pfu of Prom Δ LAT Δ 34.5-GFP in 2 μ l of MEM/FBS/antibiotics medium at the same location as the tumor implantation. This was followed by regular intraperitoneal injections of gancyclovir (7.5 mg/kg every 12 hr), started 7 days after inoculation with Prom Δ LAT Δ 34.5-GFP, and 15 continued for an additional 7 days. The second group of mice received the same treatment as the first group, except that intraperitoneal injections of saline were administered instead of gancyclovir.

The third group (control) of mice received a stereotactic inoculation containing only sterile cell culture medium in the tumor site, followed by a series of intraperitoneal injections of saline as in the second group.

20 The survival of these animals was followed until day 100 post-implantation.

In parallel time course experiments done for histopathological analysis, animals in each group were sampled and euthanised at days 10, 14, 21, and 28 post-implantation. Brains were removed, the tumors were dissected out aseptically in a dissection hood, and frozen on dry ice, and frozen sections 25 were sectioned to 10 μ m thickness in a cryostat. The sections were used for detection of GFP expression or for staining with hematoxylin & eosin (HE). To determine the expression of GFP, these sections were fixed with 4 % paraformaldehyde/PBS for 5 min, then washed with phosphate buffered saline. Sections were mounted with Immuno-Fluore Mounting Medium (ICN Biomedicals, Inc., Aurora, OH) and examined by fluorescence microscopy.

A Kaplan-Meier survival curve summarizes the results in Figure 7. All mice in the control 30 group died within 33 to 44 days post-implantation. One of 10 mice (10 %) in the first group receiving Prom Δ LAT Δ 34.5-GFP and gancyclovir treatment survived at least 100 days after tumor implantation. In the second group, receiving Prom Δ LAT Δ 34.5-GFP without gancyclovir treatment, 2 of 10 mice (20 %) survived 100 days after implantation. Statistical analysis revealed that the differences of survival time between the control group and either of the Prom Δ LAT Δ 34.5-GFP treatment groups (i.e., +/- 35 gancyclovir treatment) were statistically significant ($p=0.0191$ and $p<0.0001$ respectively). However,

in this experiment there was no significant difference in survival time between the treatment group receiving gancyclovir and the treatment group not receiving gancyclovir.

Tumors from mice in either the first or second treatment groups were relatively small in size and irregularly shaped, compared with tumors in the control group. Tumors in the control group became enlarged with time and were round, containing cells possessing high cellularity and pleomorphism. Figure 8 shows HE-stained sections (left panels) and GFP expression, as detected by fluorescence microscopy (right panels). On day 10 post-implantation, HE-stained sections of malignant tumor tissue from the right basal ganglia of mice in the control group is shown in Figure 8A (left panel); control tissue lacked GFP positivity. (Fig. 8A, right panel).

10 HE-staining of tumor tissue from mice in the second treatment group, sampled at day 10 post-implantation (i.e., day 3 post-viral inoculation), revealed necrotic tumor tissue with infiltration of inflammatory cells (Fig. 8B, left).

15 On day 14, post-implantation (day 7 post-viral inoculation), tumor tissue sections exhibited necrotic and cell-lytic features. (Fig. 8C, left). On day 21 (14 days post viral inoculation; Figure 8D, left) or day 28 (21 days post-viral inoculation; data not shown), tumor tissues resembled sections from day 14, and tumor tissue sampled from the first treatment group (+gancyclovir), on Day 21, had the same features. (Fig. 8E, left) All sections of tumors from both treatment groups showed GFP positivities. (Fig. 8B-D, right).

20 On day 100, all remaining 3 survivors were euthanised and their brains were removed for histological analysis as described above. In the brain of a survivor treated with Prom Δ LAT Δ 34.5-GFP followed by intraperitoneal saline injections, HE-stained sections showed numerous cavities in the tumor tissue accompanied by necrosis. (Fig. 9A, left panel). Tumor cells had GFP positivities. (Figure 9A, right panel)

25 In the surviving mouse that received Prom Δ LAT Δ 34.5-GFP and gancyclovir injections, tumors were relatively small in size considering the survival date of 100 days. Tumor tissue appeared necrotic or cell-lytic and was surrounded by a gliosis. (Figure 9B, left panel). The tumor cells were positive for GFP expression. (Fig. 9B, right panel).

The foregoing examples being illustrative but not an exhaustive description of the embodiments of the present invention, the following claims are presented.

CLAIMS

1. A method of selectively inhibiting the growth of a malignant cell in a mammal, comprising:

delivering to a malignant cell in a mammal an HSV-1-derived viral vector containing a DNA having a deletion in both copies of the LAT gene and a deletion in both copies of the ICP34.5 gene of HSV-1; and

causing said cell to internalize and express said vector, transcripts encoding LAT gene product and ICP34.5 gene product not being detectably produced in said malignant cell, whereby inhibition of the growth of said malignant cell results.

2. The method of Claim 1, wherein said malignant cell is derived from or is contained in a glioma, glioblastoma, oligodendrolioma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma.

3. The method of Claim 1, wherein the malignant cell is contained in a malignancy in the skull, brain, spine, thorax, lung, peritoneum, prostate, ovary, uterus, breast, stomach, liver, bowel, colon, rectum, bone, lymphatic system, or skin, of said mammal.

4. The method of Claim 1, wherein said mammal is a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

5. The method of Claim 1, wherein said HSV-1-derived vector is derived from HSV-1 strain McKrae.

6. The method of Claim 1, wherein the HSV-1-derived vector is Δ LAT Δ 34.5 or Prom Δ LAT Δ 34.5, or a derivative of either of these.

7. The method of Claim 1, wherein said HSV-1-derived vector is delivered by intrathecal injection.

8. The method of Claim 1, wherein said malignant cell is contained in a tumor, and said HSV-1-derived vector is delivered by an intratumoral injection to the malignant cell via a surgical incision.

9. The method of Claim 8, wherein the surgical incision is a craniotomy.

10. The method of Claim 8, further comprising surgically debulking the tumor containing the malignant cell before delivering said HSV-1-derived vector.

11. The method of Claim 1, wherein said HSV-1-derived vector is delivered by stereotactic inoculation.

12. The method of Claim 1, wherein said HSV-1-derived vector is delivered transvascularly.

13. The method of Claim 12, wherein transvascular delivery is by intravenous or intra-arterial injection.

14. The method of Claim 13, further comprising disrupting the blood-brain barrier before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

15. The method of Claim 14, wherein disruption of the blood-brain barrier is accomplished by administering to the mammal a hypertonic infusion.

16. The method of Claim 15, wherein the hypertonic infusion is a mannitol infusion.

17. The method of Claim 13, further comprising administering to said mammal a vasoactive agent before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

18. The method of Claim 17, wherein said vasoactive agent is bradykinin or a functional analog thereof.

19. A method of selectively inhibiting the growth of a malignant cell in a mammal, comprising:

delivering to a malignant cell in a mammal an HSV-1-derived vector containing a DNA having a deletion in both copies of the LAT gene and a deletion in both copies of the ICP34.5 gene of HSV-1, the DNA of said vector comprising at least one transcriptional unit of a LAT promoter sequence or operative fragment thereof, operatively linked to a nucleic acid sequence encoding a protein, or functional fragment thereof, toxic to said malignant cell; and

causing said cell to internalize said vector and express, from said transcriptional unit, transcript encoding said toxic protein, or functional fragment thereof, transcripts encoding LAT gene product and ICP34.5 gene product not being detectably produced in said malignant cell, whereby said toxic protein, or functional fragment thereof, is produced in said malignant cell, resulting in an inhibition of the growth of said malignant cell.

20. The method of Claim 19, wherein said malignant cell is derived from or is contained in a glioma, glioblastoma, oligodendrolioma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma.

21. The method of Claim 19, wherein the malignant cell is contained in a malignancy in the skull, brain, spine, thorax, lung, peritoneum, prostate, ovary, uterus, breast, stomach, liver, bowel, colon, rectum, bone, lymphatic system, or skin, of said mammal.

22. The method of Claim 19, wherein said mammal is a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

23. The method of Claim 19, wherein said HSV-1-derived vector is derived from HSV-1 strain McKrae.

24. The method of Claim 19, wherein the HSV-1-derived vector is Δ LAT Δ 34.5 or Prom Δ LAT Δ 34.5, or a derivative of either of these.

25. The method of Claim 19, wherein said HSV-1-derived vector is delivered by intrathecal injection.

26. The method of Claim 19, wherein said malignant cell is contained in a tumor, and said HSV-1-derived vector is delivered by an intratumoral injection to the malignant cell via a surgical incision.

27. The method of Claim 26, wherein the surgical incision is a craniotomy.

28. The method of Claim 26, further comprising surgically debulking the tumor containing the malignant cell before delivering said HSV-1-derived vector.

29. The method of Claim 19, wherein said HSV-1-derived vector is delivered by stereotactic inoculation.

30. The method of Claim 19, wherein said HSV-1-derived vector is delivered transvascularly.

31. The method of Claim 30, wherein transvascular delivery is by intravenous or intra-arterial injection.

32. The method of Claim 31, further comprising disrupting the blood-brain barrier before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

33. The method of Claim 32, wherein disruption of the blood-brain barrier is accomplished by administering to the mammal a hypertonic infusion.

34. The method of Claim 33, wherein the hypertonic infusion is a mannitol infusion.

35. The method of Claim 31, further comprising administering to said mammal a vasoactive agent before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

36. The method of Claim 35, wherein said vasoactive agent is bradykinin or a functional analog thereof.

37. The method of Claim 19, wherein said toxic protein is human interferon- γ , interleukin-2, interleukin-4, interleukin-6, interleukin-10, interleukin-12, granulocyte-macrophage colony stimulating factor, tumor necrosis factor- α , Fas ligand, human connexin-43, VP-16, or VP-22, or a functional fragment or fusion protein derived from any of these.

38. A method of selectively inhibiting the growth of a malignant cell in a mammal, comprising:

delivering to a malignant cell in a mammal an HSV-1-derived vector containing a DNA having a deletion in both copies of the LAT gene and a deletion in both copies of the ICP34.5 gene of HSV-1, the DNA of said vector comprising at least one transcriptional unit of a LAT promoter

sequence or operative fragment thereof, operatively linked to a nucleic acid sequence encoding a human interferon- γ , interleukin-2, interleukin-4, interleukin-6, interleukin-10, interleukin-12, granulocyte-macrophage colony stimulating factor, tumor necrosis factor- α , Fas ligand, human connexin-43, VP-16, or VP-22, or a functional fragment or fusion protein derived from any of these; and

causing said cell to internalize said vector and express transcript from said transcriptional unit, transcripts encoding LAT gene product and ICP34.5 gene product not being detectably produced in said malignant cell, whereby human interferon- γ , interleukin-2, interleukin-4, interleukin-6, interleukin-10, interleukin-12, granulocyte-macrophage colony stimulating factor, tumor necrosis factor- α , Fas ligand, human connexin-43, VP-16, VP-22, or a functional fragment or fusion protein derived from any of these, is produced, resulting in an inhibition of the growth of said malignant cell.

39. The method of Claim 38, wherein said malignant cell is derived from or is contained in a glioma, glioblastoma, oligodendrolioma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma.

40. The method of Claim 38, wherein the malignant cell is contained in a malignancy in the skull, brain, spine, thorax, lung, peritoneum, prostate, ovary, uterus, breast, stomach, liver, bowel, colon, rectum, bone, lymphatic system, or skin, of said mammal.

41. The method of Claim 38, wherein said mammal is a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

42. The method of Claim 38, wherein said HSV-1-derived vector is derived from HSV-1 strain McKrae.

43. The method of Claim 38, wherein the HSV-1-derived vector is Δ LAT Δ 34.5 or Prom Δ LAT Δ 34.5, or a derivative of either of these.

44. The method of Claim 38, wherein said malignant cell is contained in a tumor, and said HSV-1-derived expression vector is delivered by an intratumoral injection to the malignant cell via a surgical incision.

45. The method of Claim 44, wherein the surgical incision is a craniotomy.

46. The method of Claim 44, further comprising surgically debulking the tumor containing the malignant cell before delivering said HSV-1-derived vector.

47. The method of Claim 38, wherein said HSV-1-derived vector is delivered by stereotactic inoculation.

48. The method of Claim 38, wherein said HSV-1-derived vector is delivered transvascularly.

49. The method of Claim 48, wherein transvascular delivery is by intra-arterial injection.

50. The method of Claim 49, further comprising disrupting the blood-brain barrier before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

51. The method of Claim 50, wherein disruption of the blood-brain barrier accomplished by administering to the mammal a hypertonic infusion.

52. The method of Claim 51, wherein the hypertonic infusion is a mannitol infusion.

53. The method of Claim 49, further comprising administering to said mammal a vasoactive agent before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

54. The method of Claim 53, wherein said vasoactive agent is bradykinin or a functional analog thereof.

55. A method of selectively inhibiting the growth of a malignant cell in a mammal, comprising:

delivering to a malignant cell in a mammal an HSV-1-derived vector containing a DNA having a deletion in both copies of the LAT gene and a deletion in both copies of the ICP34.5 gene of HSV-1, said vector, having a functional HSV thymidine kinase gene;

causing said malignant cell to internalize said vector and express transcripts encoding HSV thymidine kinase, whereby HSV thymidine kinase is produced in said malignant cell, and transcripts

encoding LAT gene product and ICP34.5 gene product are not detectably produced in said malignant cell; and

administering to said mammal a dose of gancyclovir or acyclovir, whereby gancyclovir or acyclovir is converted by HSV thymidine kinase to a nucleotide analog within said malignant cell, and whereby the growth of said malignant cell is inhibited.

56. The method of Claim 55, wherein said malignant cell is derived from or is contained in a glioma, glioblastoma, oligodendrolioma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma.

57. The method of Claim 55, wherein the malignant cell is contained in a malignancy in the skull, brain, spine, thorax, lung, peritoneum, prostate, ovary, uterus, breast, stomach, liver, bowel, colon, rectum, bone, lymphatic system, or skin, of said mammal.

58. The method of Claim 55, wherein said mammal is a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

59. The method of Claim 55, wherein said HSV-1-derived vector is derived from HSV-1 strain McKrae.

60. The method of Claim 55, wherein the HSV-1-derived vector is Δ LAT Δ 34.5 or Prom Δ LAT Δ 34.5, or a derivative of either of these.

61. The method of Claim 55, wherein said malignant cell is contained in a tumor, and said HSV-1-derived vector is delivered by an intratumoral injection to the malignant cell via a surgical incision.

62. The method of Claim 61, wherein the surgical incision is a craniotomy.

63. The method of Claim 61, further comprising surgically debulking the tumor containing the malignant cell before delivering said HSV-1-derived expression vector.

64. The method of Claim 55, wherein said HSV-1-derived expression vector is delivered by stereotactic inoculation.

65. The method of Claim 55, wherein said HSV-1-derived expression vector is delivered transvascularly.

66. The method of Claim 65, wherein transvascular delivery is by intravenous or intra-arterial injection.

67. The method of Claim 66, further comprising disrupting the blood-brain barrier before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

68. The method of Claim 67, wherein disruption of the blood-brain barrier accomplished by administering to the mammal a hypertonic infusion.

69. The method of Claim 68, wherein the hypertonic infusion is a mannitol infusion.

70. The method of Claim 66, further comprising administering to said mammal a vasoactive agent before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

71. The method of Claim 70, wherein said vasoactive agent is bradykinin or a functional analog thereof.

72. A method of expressing in a mammalian cell a gene encoding a preselected protein, comprising:

delivering to a mammalian cell an HSV-1-derived vector containing a DNA having a deletion in both copies of the LAT gene and a deletion in both copies of the ICP34.5 gene of HSV-1, the DNA of said vector comprising at least one transcriptional unit of a LAT promoter sequence or operative fragment thereof, operatively linked to a nucleic acid sequence encoding a preselected protein; and

causing said cell to internalize said vector and express transcript encoding said preselected protein from said transcriptional unit, transcripts encoding LAT gene product and ICP34.5 gene product not being detectably produced in said cell, said preselected protein being produced in said cell, whereby said HSV-expressing cell produces an elevated amount of said desired protein compared to a homologous uninfected cell.

73. The method of Claim 72, wherein said cell is a malignant cell.

74. The method of Claim 72, wherein said cell is a non-malignant cell.

75. The method of Claim 72, wherein said cell is a malignant cell derived from or is contained in a glioma, glioblastoma, oligodendrolioma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma.

76. The method of Claim 72, wherein the malignant cell is derived from or is contained in a malignancy in the skull, brain, spine, thorax, lung, peritoneum, prostate, ovary, uterus, breast, stomach, liver, bowel, colon, rectum, bone, lymphatic system, or skin, of a mammal.

77. The method of Claim 72, wherein said mammalian cell is derived from or is contained in a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

78. The method of Claim 72, wherein said HSV-1-derived vector is derived from HSV-1 strain McKrae.

79. The method of Claim 72, wherein the HSV-1-derived vector is PromΔLATΔ34.5, or a derivative thereof.

80. The method of Claim 72, wherein the preselected protein is a fluorescent or light-emitting protein or a cytokine.

81. The method of Claim 80, wherein said fluorescent or light-emitting protein is a green fluorescent protein, yellow fluorescent protein, blue fluorescent protein, phycobiliprotein, luciferase, or apoaequorin.

82. The method of Claim 72, further comprising introducing said HSV-1-derived vector to a mammalian cell in vitro.

83. The method of Claim 72, further comprising introducing said HSV-1-derived vector to the mammalian cell within a mammal in vivo.

84. The method of Claim 83, wherein said mammal is a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

85. The method of Claim 83, wherein said mammalian cell is a malignant cell contained in a tumor, and said HSV-1-derived expression vector is delivered by an intratumoral injection to the malignant cell via a surgical incision.

86. The method of Claim 83, wherein the surgical incision is a craniotomy.

87. The method of Claim 85, further comprising surgically debulking the tumor containing the cell before delivering said HSV-1-derived expression vector.

88. The method of Claim 83, wherein said HSV-1-derived vector is delivered by stereotactic inoculation.

89. The method of Claim 83, wherein said HSV-1-derived vector is delivered transvascularly.

90. The method of Claim 89, wherein transvascular delivery is by intra-arterial injection.

91. The method of Claim 90, further comprising disrupting the blood-brain barrier before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

92. The method of Claim 91, wherein disruption of the blood-brain barrier is accomplished by administering to the mammal a hypertonic infusion.

93. The method of Claim 92, wherein the hypertonic infusion is a mannitol infusion.

94. The method of Claim 90, further comprising administering to said mammal a vasoactive agent before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

95. The method of Claim 94, wherein said vasoactive agent is bradykinin or a functional analog of bradykinin.

96. A method of treating a genetic defect in a mammal, comprising the method of

Claim 72.

97. A method of detecting the presence of a cell expressing HSV-1, comprising: delivering to a mammalian cell an HSV-1-derived vector containing DNA having a deletion in both copies of the LAT gene and a deletion in both copies of the ICP34.5 gene of HSV-1, the DNA of said vector comprising at least one transcriptional unit of a LAT promoter sequence or operative fragment thereof, operatively linked to a nucleic acid sequence encoding a fluorescent or light-emitting protein; and

causing said cell to internalize said vector and express transcript from said transcriptional unit, transcripts encoding LAT gene product and ICP34.5 gene product not being detectably produced in said cell, said fluorescent or light-emitting protein being produced in said cell, whereby said HSV-expressing cell is detected via luminescence or fluorescence therefrom.

98. The method of Claim 97, wherein said cell is a malignant cell.

99. The method of Claim 97, wherein said cell is a non-malignant cell.

100. The method of Claim 97, wherein said cell is a malignant cell derived from or contained in a glioma, glioblastoma, oligodendrolioma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma.

101. The method of Claim 97, wherein the cell is derived from or is contained in a malignancy in the skull, brain, spine, thorax, lung, peritoneum, prostate, ovary, uterus, breast, stomach, liver, bowel, colon, rectum, bone, lymphatic system, or skin, of a mammal.

102. The method of Claim 97, wherein said mammalian cell is derived from or is contained in a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

103. The method of Claim 97, wherein said HSV-1-derived vector is derived from HSV-1 strain McKrae.

104. The method of Claim 97, wherein the HSV-1-derived vector is Prom Δ LAT Δ 34.5, or a derivative thereof.

105. The method of Claim 97, wherein said fluorescent or light-emitting protein is a green

fluorescent protein, yellow fluorescent protein, blue fluorescent protein, phycobiliprotein, luciferase, or apoaequorin.

106. The method of Claim 97, further comprising introducing said HSV-1-derived vector to a mammalian cell in vitro.

107. The method of Claim 97, further comprising introducing said HSV-1-derived vector to the mammalian cell within a mammal in vivo.

108. The method of Claim 107, wherein said mammal is a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

109. The method of Claim 107, wherein said cell is a malignant cell contained in a tumor, and said HSV-1-derived vector is delivered by an intratumoral injection to the malignant cell via a surgical incision.

110. The method of Claim 109, wherein the surgical incision is a craniotomy.

111. The method of Claim 109, further comprising surgically debulking the tumor containing the cell before delivering said HSV-1-derived vector.

112. The method of Claim 107, wherein said HSV-1-derived vector is delivered by stereotactic inoculation.

113. The method of Claim 107, wherein said HSV-1-derived vector is delivered transvascularly.

114. The method of Claim 113, wherein transvascular delivery is by intra-arterial injection.

115. The method of Claim 114, further comprising disrupting the blood-brain barrier before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

116. The method of Claim 115, wherein disruption of the blood-brain barrier is accomplished by administering to the mammal a hypertonic infusion.

117. The method of Claim 116, wherein the hypertonic infusion is a mannitol infusion.

118. The method of Claim 114, further comprising administering to said mammal a vasoactive agent before or substantially simultaneously with intra-arterial injection of said HSV-derived vector, whereby passage of said HSV-1-derived vector across the blood-brain barrier is facilitated.

119. The method of Claim 118, wherein said vasoactive agent is bradykinin or a functional analog thereof.

120. An HSV-1-derived vector, having a functional HSV thymidine kinase gene, a deletion in both copies of the HSV-1 LAT gene, and a deletion in both copies of the HSV ICP34.5 gene, such that functional RNA transcripts encoding the LAT gene product and encoding the ICP34.5 gene product cannot be detected within a mammalian cell hosting said vector.

121. The HSV-derived vector of Claim 120, wherein said HSV-1-derived expression vector is derived from HSV-1 strain McKrae.

122. The HSV-1-derived vector of Claim 121, wherein the HSV-1-derived vector is Δ LAT Δ 34.5, Prom Δ LAT Δ 34.5, Prom Δ LAT Δ 34.5-GFP, or a derivative of any of these.

123. The HSV-1-derived vector of Claim 120, further comprising a DNA having at least one nucleic acid sequence defining a functional LAT promoter.

124. The HSV-1-derived vector of Claim 120, further comprising a DNA having at least one transcriptional unit having a functional LAT Promoter sequence or operative fragment thereof, operatively linked to nucleic acid encoding a preselected protein.

125. The HSV-1-derived vector of Claim 124, wherein the preselected protein is a fluorescent or light-emitting protein.

126. The HSV-1-derived vector of Claim 125, wherein said fluorescent or light-emitting protein is a green fluorescent protein, yellow fluorescent protein, blue fluorescent protein, phycobiliprotein, luciferase, or apoaequorin.

127. The HSV-1-derived vector of Claim 126, wherein the vector is Prom Δ LAT Δ 34.5-

GFP or a derivative thereof.

128. The HSV-1-derived vector of Claim 123, wherein the HSV-1-derived vector is Prom Δ LAT Δ 34.5, Prom Δ LAT Δ 34.5-GFP, or a derivative of either of these.

129. The HSV-1-derived vector of Claim 124, wherein the preselected protein is a protein toxic to cells expressing HSV.

130. The HSV-1-derived vector of Claim 124, wherein the preselected protein is human interferon- γ , interleukin-2, interleukin-4, interleukin-6, interleukin-10, interleukin-12, granulocyte-macrophage colony stimulating factor, tumor necrosis factor- α , Fas ligand, human connexin-43, VP-16, or VP-22, or a fusion protein derived from any of these.

131. An HSV-1-derived vector, comprising a DNA having at least one nucleic acid sequence defining a functional LAT promoter sequence or operative fragment thereof, and having a deletion in both copies of the HSV-1 LAT gene and a deletion in both copies of the HSV ICP34.5 gene, such that functional RNA transcripts encoding the LAT gene product and encoding the ICP34.5 gene product cannot be detected within a mammalian cell hosting said vector.

132. The HSV-1-derived vector of Claim 131, wherein said HSV-1-derived vector is derived from HSV-1 strain McKrae.

133. The HSV-1-derived vector of Claim 131, further comprising a functional HSV thymidine kinase gene.

134. The HSV-1-derived vector of Claim 132, wherein the HSV-1-derived vector is Prom Δ LAT Δ 34.5, Prom Δ LAT Δ 34.5-GFP, or a derivative of either of these.

135. The HSV-1-derived vector of Claim 131, further comprising a DNA having at least one transcriptional unit having a LAT Promoter sequence or operative fragment thereof, operatively linked to a nucleic acid encoding a preselected protein.

136. The HSV-1-derived vector of Claim 135, wherein the preselected protein is a fluorescent or light-emitting protein or cytokine.

137. The HSV-1-derived vector of Claim 136, wherein said fluorescent or light-emitting

protein is a green fluorescent protein, yellow fluorescent protein, blue fluorescent protein, phycobiliprotein, luciferase, or apoaequorin.

138. The HSV-1-derived vector of Claim 137, wherein the vector is Prom Δ LAT Δ 34.5-GFP or a derivative thereof.

139. The HSV-1-derived vector of Claim 131, wherein the preselected protein is a protein toxic to a cell expressing HSV.

140. The HSV-1-derived vector of Claim 131, wherein the preselected protein is human interferon- γ , interleukin-2, interleukin-4, interleukin-6, interleukin-10, interleukin-12, granulocyte-macrophage colony stimulating factor, tumor necrosis factor- α , Fas ligand, human connexin-43, VP-16, or VP-22, or a functional fragment or fusion protein derived from any of these.

141. A mammalian cell containing an HSV-1-derived vector selected from the group consisting of Δ LAT Δ 34.5, Prom Δ LAT Δ 34.5, and Prom Δ LAT Δ 34.5-GFP, or containing a derivative or unpackaged DNA of any of these.

142. The mammalian cell of Claim 141, wherein said cell is a malignant cell.

143. The mammalian cell of Claim 141, wherein said cell is a non-malignant cell.

144. The mammalian cell of Claim 141, wherein said cell is a malignant cell derived from or is contained in a glioma, glioblastoma, oligodendrogloma, astrocytoma, ependymoma, primitive neuroectodermal tumor, atypical meningioma, malignant meningioma, neuroblastoma, sarcoma, lymphoma, or carcinoma.

145. The mammalian cell of Claim 141, wherein said mammalian cell is derived from or is contained in a human, non-human primate, mouse, rat, gerbil, hamster, or rabbit.

146. A mammalian cell containing the HSV-1-derived vector of Claim 120 or unpackaged DNA thereof.

147. A mammalian cell containing the HSV-1-derived vector of Claim 121 or unpackaged DNA thereof.

148. A mammalian cell containing the HSV-1-derived vector of Claim 123 or unpackaged DNA thereof.
149. A mammalian cell containing the HSV-1-derived vector of Claim 124 or unpackaged DNA thereof.
150. A mammalian cell containing the HSV-1-derived vector of Claim 125 or unpackaged DNA thereof.
151. A mammalian cell containing the HSV-1-derived vector of Claim 126 or unpackaged DNA thereof.
152. A mammalian cell containing the HSV-1-derived vector of Claim 129 or unpackaged DNA thereof.
153. A mammalian cell containing the HSV-1-derived vector of Claim 130 or unpackaged DNA thereof.
154. A mammalian cell containing the HSV-1-derived vector of Claim 131 or unpackaged DNA thereof.
155. A mammalian cell containing the HSV-1-derived vector of Claim 132 or unpackaged DNA thereof.
156. A mammalian cell containing the HSV-1-derived vector of Claim 133 or unpackaged DNA thereof.
157. A mammalian cell containing the HSV-1-derived vector of Claim 135 or unpackaged DNA thereof.
158. A mammalian cell containing the HSV-1-derived vector of Claim 136 or unpackaged DNA thereof.
159. A mammalian cell containing the HSV-1-derived vector of Claim 137 or unpackaged DNA thereof.

160. A mammalian cell containing the HSV-1-derived vector of Claim 139 or unpackaged DNA thereof.

161. A mammalian cell containing the HSV-1-derived vector of Claim 140 or unpackaged DNA thereof.

162. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing an HSV-1-derived vector selected from the group consisting of Δ LAT Δ 34.5, Prom Δ LAT Δ 34.5, and Prom Δ LAT Δ 34.5-GFP, or containing a derivative of any of these.

163. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 120.

164. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 121.

165. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 122.

166. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 124.

167. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 125.

168. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 126.

169. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 129.

170. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 130.

171. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 131.

172. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 132.

173. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 133.

174. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 135.

175. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 136.

176. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 137.

177. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 139.

178. A kit for expressing in a mammalian cell a gene encoding a preselected protein, containing the HSV-1-derived vector of Claim 140.

179. The method of Claim 1, wherein a bispecific antibody is used to cause said cell to internalize said vector.

180. The method of Claim 19, wherein a bispecific antibody is used to cause said cell to internalize said vector.

181. The method of Claim 38, wherein a bispecific antibody is used to cause said cell to internalize said vector.

182. The method of Claim 55, wherein a bispecific antibody is used to cause said cell to internalize said vector.

183. The method of Claim 72, wherein a bispecific antibody is used to cause said cell to internalize said vector.

184. The method of Claim 97, wherein a bispecific antibody is used to cause said cell to internalize said vector.

1/9




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2

3/9




FIG. 3A




FIG. 3B

4/9




FIG. 3C

5/9

FIG. 4A




FIG. 4B




FIG. 4C






FIG. 5A

FIG. 5B

FIG. 5C

7/9




FIG. 6




FIG. 7

FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D





FIG. 8E



9/9



FIG. 9A




FIG. 9B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/11031

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/869 C12N5/06 A61K39/245 A61K48/00 A61K35/76
C07K14/035 C12N15/38 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A61K C07K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^a	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 92 13943 A (SMITHKLINE BEECHAM BIOLOG) 20 August 1992 (1992-08-20)</p> <p>claim 3 page 3, line 2,3 page 17, line 18-20</p> <p>---</p> <p>-/-</p>	<p>120-123, 131-134, 141-148, 154-156, 162-165, 171-173</p>

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

^a Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

17 July 2000

Date of mailing of the international search report

04.09.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herrmann, K

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/11031

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>MCGEOCH D J ET AL: "COMPARATIVE SEQUENCE ANALYSIS OF THE LONG REPEAT REGIONS AND ADJOINING PARTS OF THE LONG UNIQUE REGIONS IN THE GENOMES OF HERPES SIMPLEX VIRUSES TYPES 1 AND 2" JOURNAL OF GENERAL VIROLOGY 1991, vol. 72, no. 12, 1991, pages 3057-3076, XP000406755 ISSN: 0022-1317 abstract</p> <p style="text-align: center;">---</p>	120-123, 131-134, 141-148, 154-156, 162-165, 171-173
Y	<p>PERNG GUEY-CHUEN ET AL: "High-dose ocular infection with a herpes simplex virus type 1 ICP34.5 deletion mutant produces no corneal disease or neurovirulence yet results in wild-type levels of spontaneous reactivation." JOURNAL OF VIROLOGY 1996, vol. 70, no. 5, 1996, pages 2883-2893, XP002142268 ISSN: 0022-538X page 2883, paragraph 2</p> <p style="text-align: center;">---</p>	1-184
Y	<p>PERNG GUEY-CHUEN ET AL: "The latency-associated transcript gene of herpes simplex virus type 1 (HSV-1) is required for efficient in vivo spontaneous reactivation of HSV-1 from latency." JOURNAL OF VIROLOGY 1994, vol. 68, no. 12, 1994, pages 8045-8055, XP000916490 ISSN: 0022-538X abstract</p> <p style="text-align: center;">---</p>	1-184
Y	<p>KRAMM CHRISTOF M ET AL: "Gene therapy for brain tumors." BRAIN PATHOLOGY 1995, vol. 5, no. 4, 1995, pages 345-381, XP000916435 ISSN: 1015-6305 the whole document</p> <p style="text-align: center;">-----</p>	1-184

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/11031

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 1-96 and 179-184 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Although claims 97-119 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/11031

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 9213943	A 20-08-1992	AT 181958	T	15-07-1999
		AU 657731	B	23-03-1995
		AU 1182992	A	07-09-1992
		CA 2101639	A	01-08-1992
		DE 69229543	D	12-08-1999
		DE 69229543	T	27-01-2000
		EP 0571410	A	01-12-1993
		ES 2136079	T	16-11-1999
		GR 3031358	T	31-01-2000
		JP 6507066	T	11-08-1994
		NZ 241476	A	23-12-1993
		NZ 245842	A	23-12-1993
		US 6040169	A	21-03-2000
		ZA 9200651	A	24-02-1993